Projected Annual Recreational Harvest by Region for Selected Management Options

Using 2004 and 2005 CRFS catch estimates

(2005 data includes projections for Dec)

Baseline Season: Each table provides expected future catch using the 2005 management structure (what was in place at end of 2005) as the baseline season with the following changes:

no changes

	Region (see bo	ttom of page fo	r definitions)				
Species Group Name	1 (N)	2 (N-C)	3 (S-C Mont)	4 (S-C MB)	5 (S)	Grand Total (mt)*	2006 HT**
Black Rockfish	114.8	55.0	10.8	0.1	=	180.7	171
Bocaccio - South of 40° 30'		0.2	1.8	1.8	41.8	45.6	60
Cabezon	5.4	22.8	2.3	1.5	4.9	36.8	42.1
California Sheephead	-	1	-	-	30.6	30.6	61.5
Canary Rockfish	0.4	3.1	0.1	0.4	0.3	4.4	9.3
Cowcod - South of 40° 30'		-	-	-	0.2	0.2	0.4
Deeper Nearshore Rockfish - South of 40° 10'		151.3	89.9	55.5	50.3	346.9	245
Greenlings	1.6	4.3	0.9	0.1	0.0	6.9	15.5
Lingcod	50.6	135.3	26.3	17.2	24.7	254.2	422
Minor Nearshore Rockfish - North of 40° 10'	18.3					18.3	11
Scorpionfish		-	-	-	30.0	30.0	137
Shallow Nearshore Rockfish - South of 40° 10'		20.7	13.3	9.2	5.2	48.4	66
Widow Rockfish	0.0	2.0	0.1	0.0	3.9	6.0	7
Yelloweye Rockfish	0.4	0.3	0.0	0.0	0.0	0.8	3.7
Minor Nearshore Rockfish - South of 40° 10'		171.9	103.2	64.7	55.4	395.3	311

Note:

"0.0" = value between 0 and 0.05

" - " = value of 0.0

* does not include shore-based catch

** HT = Harvest Target

Regions:

(1) N = North Region: North of $40^{\circ}10'$ N lat to CA/OR border

(2) N-C = North-Central Region: South of 40°10' N lat to 37°11' N lat (Pigeon Pt.)

(3) S-C Mon = South-Central Monterey Region: South of 37°11' N lat (Pigeon Pt.) to 36° N lat (Lopez Pt.)

(4) S-C MB = South-Central Morro Bay Region: South of 36° N lat (Lopez Pt.) to 34°27' N lat (Pt. Conception)

(5) S = South Region: South of 34°27' N lat (Pt. Conception) to CA/Mexico Border

Prepared for: Groundfish Task Force meeting, 1/11/06

Prepared By: K. Lynn, CDFG

Draft Version 1/11/06; Revision 1/26/06

California Department of Fish and Game Projected Annual Recreational Harvest by Region for Selected Management Options

Using 2004 and 2005 CRFS catch estimates

(2005 data includes projections for Dec)

Baseline Season: Each table provides expected future catch using the 2005 management structure (what was in place at end of 2005) as the baseline season with the following changes:

S: 0-60 fm March (changed from 30-60 fm)

	Region (see bo	ttom of page fo	r definitions)				
Species Group Name	1 (N)	2 (N-C)	3 (S-C Mont)	4 (S-C MB)	5 (S)	Grand Total (mt)*	2006 HT**
Black Rockfish	114.8	55.0	10.8	0.1	-	180.7	171
Bocaccio - South of 40° 30'		0.2	1.8	1.8	41.3	45.2	60
Cabezon	5.4	22.8	2.3	1.5	5.4	37.3	42.1
California Sheephead	-	-	-	-	32.7	32.7	61.5
Canary Rockfish	0.4	3.1	0.1	0.4	0.3	4.4	9.3
Cowcod - South of 40° 30'		-	-	-	0.2	0.2	0.4
Deeper Nearshore Rockfish - South of 40° 10'		151.3	89.9	55.5	55.2	351.9	245
Greenlings	1.6	4.3	0.9	0.1	0.0	6.9	15.5
Lingcod	50.6	135.3	26.3	17.2	24.7	254.2	422
Minor Nearshore Rockfish - North of 40° 10'	18.3					18.3	11
Scorpionfish		-	-	-	30.2	30.2	137
Shallow Nearshore Rockfish - South of 40° 10'		20.7	13.3	9.2	5.7	49.0	66
Widow Rockfish	0.0	2.0	0.1	0.0	3.6	5.7	7
Yelloweye Rockfish	0.4	0.3	0.0	0.0	0.0	0.8	3.7
Minor Nearshore Rockfish - South of 40° 10'		171.9	103.2	64.7	61.0	400.8	311

Note:

"0.0" = value between 0 and 0.05

" - " = value of 0.0

Regions:

(1) N = North Region: North of 40°10' N lat to CA/OR border

(2) N-C = North-Central Region: South of 40°10' N lat to 37°11' N lat (Pigeon Pt.)

(3) S-C Mon = South-Central Monterey Region: South of 37°11' N lat (Pigeon Pt.) to 36° N lat (Lopez Pt.)

(4) S-C MB = South-Central Morro Bay Region: South of 36° N lat (Lopez Pt.) to 34°27' N lat (Pt. Conception)

(5) S = South Region: South of 34°27' N lat (Pt. Conception) to CA/Mexico Border

* does not include shore-based catch

** HT = Harvest Target

Prepared for: Groundfish Task Force meeting, 1/11/06

Prepared By: K. Lynn, CDFG

Draft Version 1/11/06; Revision 1/26/06

Projected Annual Recreational Harvest by Region for Selected Management Options

Using 2004 and 2005 CRFS catch estimates

(2005 data includes projections for Dec)

Baseline Season: Each table provides expected future catch using the 2005 management structure (what was in place at end of 2005) as the baseline season with the following changes:

S: 0-60 fm March (changed from 30-60 fm)
N-C: 0-30 fm season (changed from 0-20 fm)
S-C Mont: 0-30 fm season (changed from 0-20 fm)

S-C MB: add April (changed from May-Sep, still at 0-40 fm)

	Region (see bo	ttom of page fo	r definitions)				
Species Group Name	1 (N)	2 (N-C)	3 (S-C Mont)	4 (S-C MB)	5 (S)	Grand Total (mt)*	2006 HT**
Black Rockfish	114.8	51.3	10.0	0.1	-	176.2	171
Bocaccio - South of 40° 30'		0.2	5.6	2.2	41.3	49.4	60
Cabezon	5.4	20.9	2.1	1.6	5.4	35.4	42.1
California Sheephead	-	-	-	-	32.7	32.7	61.5
Canary Rockfish	0.4	4.1	0.3	0.5	0.3	5.6	9.3
Cowcod - South of 40° 30'		-	-	-	0.2	0.2	0.4
Deeper Nearshore Rockfish - South of 40° 10'		153.9	104.2	59.5	55.2	372.7	245
Greenlings	1.6	4.5	0.8	0.1	0.0	7.0	15.5
Lingcod	50.6	135.2	27.8	19.0	24.7	257.1	422
Minor Nearshore Rockfish - North of 40° 10'	18.3					18.3	11
Scorpionfish		-	-	-	30.2	30.2	137
Shallow Nearshore Rockfish - South of 40° 10'		19.7	13.2	9.8	5.7	48.5	66
Widow Rockfish	0.0	2.3	0.2	0.1	3.6	6.1	7
Yelloweye Rockfish	0.4	0.4	0.0	0.0	0.0	0.8	3.7
Minor Nearshore Rockfish - South of 40° 10'		173.6	117.3	69.3	61.0	421.2	311

Note:

Regions:

(1) N = North Region: North of 40°10′ N lat to CA/OR border

(2) N-C = North-Central Region: South of 40°10' N lat to 37°11' N lat (Pigeon Pt.)

(3) S-C Mon = South-Central Monterey Region: South of 37°11' N lat (Pigeon Pt.) to 36° N lat (Lopez Pt.)

(4) S-C MB = South-Central Morro Bay Region: South of 36° N lat (Lopez Pt.) to 34°27' N lat (Pt. Conception)

(5) S = South Region: South of 34°27' N lat (Pt. Conception) to CA/Mexico Border

Prepared for: Groundfish Task Force meeting, 1/11/06

Prepared By: K. Lynn, CDFG

Draft Version 1/11/06; Revision 1/26/06

* does not include shore-based catch

** HT = Harvest Target

[&]quot;0.0" = value between 0 and 0.05

[&]quot; - " = value of 0.0

Projected Annual Recreational Harvest by Region for Selected Management Options

Using 2004 and 2005 CRFS catch estimates

(2005 data includes projections for Dec)

Baseline Season: Each table provides expected future catch using the 2005 management structure (what was in place at end of 2005) as the baseline season with the following changes:

S: 0-60 fm March (changed from 30-60 fm)
N-C: 0-30 fm season (changed from 0-20 fm)
S-C Mont: 0-30 fm season (changed from 0-20 fm)

S-C MB: add October (changed from May-Sep, still at 0-40 fm)

	Region (see bo	ttom of page fo	r definitions)				
Species Group Name	1 (N)	2 (N-C)	2 (N-C) 3 (S-C Mont)		5 (S)	Grand Total (mt)*	2006 HT**
Black Rockfish	114.8	51.3	10.0	0.1	-	176.3	171
Bocaccio - South of 40° 30'		0.2	5.6	2.3	41.3	49.6	60
Cabezon	5.4	20.9	2.1	1.7	5.4	35.5	42.1
California Sheephead	-	-	-	-	32.7	32.7	61.5
Canary Rockfish	0.4	4.1	0.3	0.5	0.3	5.6	9.3
Cowcod - South of 40° 30'		-	-	-	0.2	0.2	0.4
Deeper Nearshore Rockfish - South of 40° 10'		153.9	104.2	70.7	55.2	384.0	245
Greenlings	1.6	4.5	0.8	0.1	0.0	7.0	15.5
Lingcod	50.6	135.2	27.8	21.5	24.7	259.7	422
Minor Nearshore Rockfish - North of 40° 10'	18.3					18.3	11
Scorpionfish		-	-	-	30.2	30.2	137
Shallow Nearshore Rockfish - South of 40° 10'		19.7	13.2	11.9	5.7	50.5	66
Widow Rockfish	0.0	2.3	0.2	0.1	3.6	6.1	7
Yelloweye Rockfish	0.4	0.4	0.0	0.0	0.0	0.8	3.7
Minor Nearshore Rockfish - South of 40° 10'		173.6	117.3	82.6	61.0	434.5	311

Note:

Regions:

(1) N = North Region: North of 40°10' N lat to CA/OR border

(2) N-C = North-Central Region: South of 40°10' N lat to 37°11' N lat (Pigeon Pt.)

(3) S-C Mon = South-Central Monterey Region: South of 37°11' N lat (Pigeon Pt.) to 36° N lat (Lopez Pt.)

(4) S-C MB = South-Central Morro Bay Region: South of 36° N lat (Lopez Pt.) to 34°27' N lat (Pt. Conception)

(5) S = South Region: South of 34°27' N lat (Pt. Conception) to CA/Mexico Border

Prepared for: Groundfish Task Force meeting, 1/11/06

Prepared By: K. Lynn, CDFG

Draft Version 1/11/06; Revision 1/26/06

* does not include shore-based catch

** HT = Harvest Target

[&]quot;0.0" = value between 0 and 0.05

[&]quot; - " = value of 0.0

Projected Annual Recreational Harvest by Region for Selected Management Options

Using 2004 and 2005 CRFS catch estimates

(2005 data includes projections for Dec)

Baseline Season: Each table provides expected future catch using the 2005 management structure (what was in place at end of 2005) as the baseline season with the following changes:

S: 0-60 fm March (changed from 30-60 fm)
N-C: 0-30 fm season (changed from 0-20 fm)

S-C Mont: 0-30 fm season (changed from 0-20 fm)

S-C MB: add April and October (changed from May-Sep, still at 0-40 fm)

	Region (see bo	ttom of page fo	r definitions)				
Species Group Name	1 (N)	2 (N-C)	3 (S-C Mont)	4 (S-C MB)	5 (S)	Grand Total (mt)*	2006 HT**
Black Rockfish	114.8	51.3	10.0	0.2	-	176.3	171
Bocaccio - South of 40° 30'		0.2	5.6	2.7	41.3	49.9	60
Cabezon	5.4	20.9	2.1	1.9	5.4	35.7	42.1
California Sheephead	-	-	-	-	32.7	32.7	61.5
Canary Rockfish	0.4	4.1	0.3	0.6	0.3	5.7	9.3
Cowcod - South of 40° 30'		-	-	-	0.2	0.2	0.4
Deeper Nearshore Rockfish - South of 40° 10'		153.9	104.2	74.7	55.2	388.0	245
Greenlings	1.6	4.5	0.8	0.1	0.0	7.0	15.5
Lingcod	50.6	135.2	27.8	23.3	24.7	261.4	422
Minor Nearshore Rockfish - North of 40° 10'	18.3					18.3	11
Scorpionfish		-	-	-	30.2	30.2	137
Shallow Nearshore Rockfish - South of 40° 10'		19.7	13.2	12.5	5.7	51.1	66
Widow Rockfish	0.0	2.3	0.2	0.1	3.6	6.1	7
Yelloweye Rockfish	0.4	0.4	0.0	0.0	0.0	0.8	3.7
Minor Nearshore Rockfish - South of 40° 10'		173.6	117.3	87.2	61.0	439.1	311

Note:

Regions:

(1) N = North Region: North of 40°10' N lat to CA/OR border

(2) N-C = North-Central Region: South of 40°10' N lat to 37°11' N lat (Pigeon Pt.)

(3) S-C Mon = South-Central Monterey Region: South of 37°11' N lat (Pigeon Pt.) to 36° N lat (Lopez Pt.)

(4) S-C MB = South-Central Morro Bay Region: South of 36° N lat (Lopez Pt.) to 34°27' N lat (Pt. Conception)

(5) S = South Region: South of 34°27' N lat (Pt. Conception) to CA/Mexico Border

Prepared for: Groundfish Task Force meeting, 1/11/06

Prepared By: K. Lynn, CDFG

Draft Version 1/11/06; Revision 1/26/06

* does not include shore-based catch

** HT = Harvest Target

[&]quot;0.0" = value between 0 and 0.05

[&]quot; - " = value of 0.0

California Recreational Regulations by Region in 2005 (Final)

Key:

	Allowed in all waters
20	Depth closed > 20fm
30	Depth closed > 30fm
40	Depth closed > 40fm
60	Depth closed > 60fm
20-40	Depth open between 20-40fm
30-60	Depth open between 30-60fm
	Closed

North Coast CA/OR Border to 40° 10' N Lat

Species	Jan	Feb	Mar	Apr	May	Jun	July	Aug	Sep	Oct	Nov	Dec
Nearshore rockfish					30	30	30	30	30	30	30	30
Black rockfish ²					30	30	30	30	30	30	30	30
California sheephead					30	30	30	30	30	30	30	30
Cabezon					30	30	30	30	30	30	30	
Greenlings (rock, kelp)					30	30	30	30	30	30	30	30
Ocean Whitefish					30	30	30	30	30	30	30	30
Shelf rockfish					30	30	30	30	30	30	30	30
Lingcod					30	30	30	30	30	30	30	

North-Central Coast

40° 10' N lat to Lopez Point (36° 00' N lat)

Species	Jan	Feb	Mar	Apr	May	Jun	July	Aug	Sep	Oct	Nov	Dec
Nearshore rockfish							20	20	20	20	20	20
California scorpionfish							20	20	20	20	20	20
California sheephead							20	20	20	20	20	20
Cabezon							20	20	20	20	20	
Greenlings (rock, kelp)							20	20	20	20	20	20
Ocean whitefish							20	20	20	20	20	20
Shelf rockfish							20	20	20	20	20	20
Lingcod							20	20	20	20	20	
Sanddabs												

South-Central Coast

Lopez Point (36° 00' N lat) to Pt. Conception (34° 27' N lat)

Species	Jan	Feb	Mar	Apr	May	Jun	July	Aug	Sep	Oct	Nov	Dec
Nearshore rockfish					40	40	40	40	40			
California scorpionfish					40	40	40	40	40			
California sheephead					40	40	40	40	40			
Cabezon					40	40	40	40	40			
Greenlings					40	40	40	40	40			
Ocean Whitefish					40	40	40	40	40			
Shelf rockfish					40	40	40	40	40			
Lingcod					40	40	40	40	40			
Sanddabs												

South Coast

Pt. Conception (34° 27' N lat) to US/Mexico Border

Species	Jan	Feb	Mar	Apr	May	Jun	July	Aug	Sep	Oct	Nov	Dec
Nearshore rockfish			30-60	60	60	60	60	60	30	30	60	60
California scorpionfish										30	60	60
California sheephead			30-60	60	60	60	60	60	30	30	60	60
Cabezon			30-60	60	60	60	60	60	30	30	60	
Greenlings			30-60	60	60	60	60	60	30	30	60	60
Ocean Whitefish			30-60	60	60	60	60	60	30	30	60	60
Shelf rockfish			30-60	60	60	60	60	60	30	30	60	60
Lingcod				60	60	60	60	60	30	30	60	
Sanddabs												